

Short communication

Evidence for pre-Variscan deformation in the Lys-Caillaouas area, Central Pyrenees, France

S.W.J. den Brok

Institute of Earth Sciences, State University of Utrecht, P.O. Box 80.021, 3508 TA Utrecht, The Netherlands

Received 4 October 1988; accepted in revised form 11 March 1989

The eastern Pyrenees are reported to show evidence of an angular unconformity between the Cambrian metasedimentary series of Canaveilles and Jujols and the Upper Ordovician Rabassa conglomerate (Llopis Lladó, 1965; Santanach Prat, 1972, 1974; Laumonier & Guitard, 1986; Laumonier, 1987, *in press*). Bedding within the Cambrian rocks trends obliquely to the lower boundary of the conglomerate (LCB). In addition, Llopis Lladó (1965) and Santanach Prat (1972, 1974) mentioned an abrupt change in orientation of δ-lineations at the LCB, i.e. a difference in orientation of intersection lineations of sedimentary bedding on the Variscan main phase schistosity. The schistosity shows no change in orientation and this implies that

bedding must have been oblique to the LCB at the time of deposition of the conglomerate.

In contrast herewith from the Central Pyrenees no observations have been reported thus far that point to an angular unconformity within the Cambro-Ordovician metasediments and the upper Ordovician conglomerate is considered to there conformably overlie the older Cambro-Ordovician rocks (Zwart, 1979; Verhoef et al., 1984; André, 1985; De Bresser et al., 1986; Speksnijder, 1987).*

However, recent observations indicate other-

* Except Alonso J.L. and Garcia Sansguido J. who describe such an unconformity in the Bosost area (*unpublished contribution to the 'Synthèse des Pyrénées*, B.R.G.M. ed., *in prep*).




Fig. 1. Geological sketch-map of Lys Caillaouas area (after Zwart 1979) showing location of the studied area.




Fig. 2. Simplified geological map of the studied area showing δ -lineations within the Camboro-Ordovician rocks of the central Lys-Caillaouas area. Each arrow indicates one measurement. Numbers indicate plunge. Large numbers, 1 to 6, mark sub areas with relatively constant orientation of main phase cleavage and/or δ -linacions. Legend: I = Silurian & Devonian; II = Upper Ordovician mica schists and quartzitic marbles; III = Upper Ordovician conglomerate; IV = pre-Upper Ordovician mica schists, quartzites and quartzitic marbles; V = porphyritic biotite granite; VI = boundary between subareas with relative constant orientation of δ -lineations and/or main phase cleavage (see Fig. 4).

wise and from the distribution of δ -lineations within the Cambro-Ordovician metasediments in the Central Lys-Caillaouas area (Central Pyrenees; see Figs. 1, 2 & 3) it now appears that: (i) the orientation of δ -lineations above the LCB, i.e. in the Upper Ordovician, is reasonably constant and plunges east with a mean value of about 15 degrees (see also Fig. 4, see page 380); (ii) there is an abrupt change in orientation across the LCB while the main phase cleavage has a constant orientation; (iii) the orientation of δ -lineations below the LCB, i.e. in the pre-Upper Ordovician, shows a considerable spread.

Conclusion

Within the Central Pyrenees the spread of δ -lineations below the LCB together with a constant orientation of the main schistosity indicates a deformational event prior to the main phase deformation. Above the LCB no significant spread of δ -lineations is found. There is an abrupt change in orientation of δ -lineations across the LCB. Pre-main phase deformation therefore only affected rocks below the LCB. The conglomerate is of Upper Ordovician age. Consequently, as in the eastern Pyrenees, in the Central Pyrenees deposition of the conglomerate must have taken place after a pre-Variscan deformational event.




Fig. 3. Cross sections through the studied area showing main phase cleavage and folding of the lower conglomerate boundary. Both folding and cleavage are of Variscan age. Legend: 1 = Upper Ordovician; 2 = pre-Upper Ordovician; 3 = porphyritic biotite granite. See figure 2 for location of sections.

Acknowledgements

Financial support from the Molengraaff Fund is gratefully acknowledged.

References

- André, J.P. 1985 Contribution à l'étude des terrains Paléozoïques de la vallée du Lys, zone axiale des Pyrénées – Bull. Soc. Geol. Fr. 1(VIII): 515–521
- De Bresser, J.H.P., F.J.M. Majoor & M. Ploegsma 1986 New insights in the structural and metamorphic history of the western Lys Caillaus massif (Central Pyrenees, France) – Geol. Mijnbouw 65: 177–187
- Laumonier, B. 1987 Révision de la lithostratigraphie du Paléozoïque inférieur des Aspres, à l'est du massif du Canigou; conséquences stratigraphiques et structurales – C.R. Acad. Sci. Paris, 305(II): 611–614
- Laumonier, B. (in press) Les groupes de Canaveilles et de Jujols ('Paléozoïque inférieur') des Pyrénées orientales; arguments en faveur de l'âge essentiellement cambrien de ces séries – Hercynica.
- Laumonier, B. & G. Guitard 1986 Le Paléozoïque inférieur de la moitié orientale de la zone axiale des Pyrénées, essai de synthèse – C.R. Acad. Sci. Paris, 302(II): 473–478
- Llopis Lladó, N. 1965 Sur le Paléozoïque inférieur de l'Andorre – Bull. Soc. Geol. Fr. 7(II): 652–659
- Santanach Prat, P.F. 1972 Sobre una discordancia en el Paleozoico inferior de los Pirineos orientales – Acta Geol. Hisp. VII-5: 129–132
- Santanach Prat, P.F. 1974 Estudi tectònic del Paleozoic Inferior del Pirineu entre la Cerdanya i el riu Ter – Fundació Salvador Vives Casajuana, Barcelona: 133 pp
- Speksnijder, A. 1987 The detection and significance of early deformation in the southern Variscan Pyrenees, Spain; implications for regional Paleozoic structural evolution – Geol. Rundsch. 76(2): 451–476
- Verhoef, P.N.W., R.L.M. Vissers & H.J. Zwart 1984 A new interpretation of the structural and metamorphic history of the western Aston massif (Central Pyrenees, France) – Geol. Mijnbouw 63: 399–410
- Zwart, H.J. 1979 The geology of the central Pyrenees – Leidse Geol. Meded. 50(1): 1–74




Fig. 4. Stereographic equal area projections per subarea showing poles on sedimentary bedding, poles on main phase cleavage and δ -lineations. Subareas are indicated in Fig. 2.